PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



(51) International Patent Classification ⁶ :		(11) International Publication Number: WO 99/47171
A61K 45/06	A1	(43) International Publication Date: 23 September 1999 (23.09.99)
(21) International Application Number: PCT/GE (22) International Filing Date: 16 March 1999 ((30) Priority Data: 9805557.7 16 March 1998 (16.03.98) (71) Applicant (for all designated States except US): SHARP & DOHME LIMITED [GB/GB]; Hertf Hoddesdon, Hertfordshire EN11 9BU (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): DAWSON, Gerar [GB/GB]; Terlings Park, Eastwick Road, Harle CM20 2QR (GB). (74) Agent: HORGAN, James; Merck & Co., Inc., Europ Dept., Terlings Park, Eastwick Road, Harlow, Ed 2QR (GB).	MERO ord Ro d, Raph ow, Es	BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GE GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, IP, KE, KC KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MI RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MI NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt amendments.

The present invention relates to a combination of a muscarinic agonist and an inverse agonist of the GABAA α_5 receptor subtype, and the use of the combination in treating neurodegenerative conditions such as Alzheimer's Disease.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL AM AT AU AZ BA BB BE BF BG BJ BR BY CA CF CG CH CI CM CN CU CZ DE DK	Albania Armenia Australia Australia Azerbaijan Bosnia and Herzegovina Barbados Belgium Burkina Faso Bulgaria Benin Brazil Belarus Canada Central African Republic Congo Switzerland Côte d'Ivoire Cameroon China Cuba Czech Republic Germany Demmark	ES FI FR GA GB GC GH GN GR HU IE IL IS IT JP KE KG KP KR LC LL LK	Spaim Finland France Gabon United Kingdom Georgia Ghana Guinea Greece Hungary Ireland Israel Iceland Italy Japan Kenya Kyrgyzstan Democratic People's Republic of Korea Republic of Korea Kazakstan Saint Lucia Liechtenstein Sri Lanka	LS LT LU LV MC MD MG MK ML MN MR MW MX NE NL NO NZ PL PT RO RU SD SE	Lesotho Lithuania Luxembourg Latvia Monaco Republic of Moldova Madagascar The former Yugoslav Republic of Macedonia Mali Mongolia Mauritania Malawi Mexico Niger Netherlands Norway New Zealand Poland Portugal Romania Russian Federation Sudan Sweden	SI SK SN SZ TD TG TJ TM TR TT UA UG US UZ VN YU ZW	Slovenia Slovakia Senegal Swaziland Chad Togo Tajikistan Turkmenistan Turkey Trinidad and Tobago Ukraine Uganda United States of America Uzbekistan Viet Nam Yugoslavia Zimbabwe
EE	Estonia	LR	Liberia	SG	Singapore		

WO 99/47171 PCT/GB99/00801

COMBINATION OF A GABA-A ALPHA 5 INVERSE AGONIST AND A MUSCARINIC AGONIST

The present invention relates to a combination of an muscarinic agonist and an inverse agonist of the GABAA α_5 receptor subtype, and the use of the combination in treating neurodegenerative conditions such as Alzheimer's Disease.

5

10

15

20

25

30

Alzheimer's Disease is a poorly understood neurodegenerative condition mainly affecting the elderly but also younger people who are generally genetically predispositioned to it.

One postulated method of treatment comprises the administration of muscarinic agonists which act on the cholinergic system. However this method suffers from the disadvantages that these compounds induce a range of side-effects including diarrhoea, salivation and nausea.

The present invention provides a new and surprisingly effective synergistic combination of an muscarinic agonist and an inverse agonist of the GABAA α_5 receptor subtype for separate, sequential or simultaneous administration.

The present invention provides a greater than expected improvement in the condition of subjects suffering from a neurodegenerative with an associated cognitive deficit, such as Alzheimer's Disease or Parkinson's disease, or from a cognitive deficit which may arise from a normal process such as aging or from an abnormal process such as injury, than would be expected from administration of the active ingredients alone. Further, the combination allows for a lower overall dose of each of the active ingredients to be administered thus reducing side effects and decreasing any reduction in the effectiveness of each of the active ingredients over time.

Muscarinic agonists which may be used include any which are known to the skilled person. Examples are methacholine and its chloride, carbachol, bethanechol, arecholine, pilocarpine, muscarine, McN-A-343,

WO 99/47171 PCT/GB99/00801

oxotremorine, milameline, xanomeline, cis-methyldioxalane, pirenzepine, gallamine, SB 202026, AF102B, AFDX 116 and RS-86.

Any inverse agonist of the GABA_A α_5 receptor subtype may be used which fulfills the criteria of WO-A-9625948. The inverse agonist may be either binding selective for the α_5 subtype or functionally selective, or both. Thus the inverse agonist is preferably an antagonist, or has insignificant agonist or inverse agonist properties at the other GABA_A α receptor subtypes when measured in oocytes as described in WO-A-9625948.

Thus the inverse agonist preferably has a functional efficacy at the α_5 receptor subunit of less than -20% and functional efficacies at the α_1 , α_2 , and α_3 receptor subunits of between -20 and +20%. By functional efficacy is meant the percentage modulation of the EC₂₀ response produced by GABA, upon coadministration of the inverse agonist, in oocytes expressing GABA_A receptor channels containing the α receptor subunit under test. Details of this measurement are given in WO-A-9625948.

The inverse agonist preferably binds selectively to GABA_A receptors containing the α_5 subunit 10, 25 and particularly 50 times compared to GABA_A receptors subunits containing the α_1 , α_2 or α_3 subunits. Preferably this binding selectivity is shown over all these subunits.

A preferred class of inverse agonists, which are disclosed in WO-A-9850385, are of formula I:

$$\begin{array}{c|c}
R^{1} & N-N \\
N & N
\end{array}$$

$$\begin{array}{c|c}
R^{2} & N-N \\
N & N
\end{array}$$

$$\begin{array}{c|c}
L-Y-X
\end{array}$$

25

5

10

15

20

wherein:

R¹ is hydrogen, halogen or CN or a group C₁.4alkyl, C₂.4alkenyl, C₂.4alkynyl, C₁.4alkoxy, C₂.4alkenyloxy or C₂.4alkynyloxy, each of which groups is unsubstituted or substituted with one or two halogen atoms or with a pyridyl or phenyl ring each of which rings may be unsubstituted or independently substituted by one or two halogen atoms or nitro, cyano, amino, methyl or CF₃ groups;

R² is hydrogen, halogen or CN or a group C_{1.4}alkyl, C_{2.4}alkenyl, C_{2.4}alkynyl, C_{1.4}alkoxy, C_{2.4}alkenyloxy or C_{2.4}alkynyloxy each of which groups is unsubstituted or substituted with one or two halogen atoms;

5

10

15

20

25

30

L is O, S or NRⁿ where Rⁿ is H, C₁₋₆alkyl or C₃₋₆cycloalkyl;

X is a 5-membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently chosen from oxygen, nitrogen and sulphur, at most one of the heteroatoms being oxygen or sulphur, or a 6-membered heteroaromatic ring containing 1, 2 or 3 nitrogen atoms, the 5- or 6-membered heteroaromatic ring being optionally fused to a benzene ring and the heteroaromatic ring being optionally substituted by Rx and/or Ry and/or Rz, where Rx is halogen, R3, OR3, OCOR3, NR4R5, NR4COR5, tri(C₁₋₆alkyl)silylC₁₋₆alkoxyC₁₋₄alkyl, CN or R⁹, R^y is halogen, R³, OR³, OCOR3, NR4R5, NR4COR5 or CN and R^z is R^3 , OR3 or OCOR3, where R^3 is $C_{1\text{-}6}$ alkyl, $C_{2\text{-}6}$ alkenyl, $C_{2\text{-}6}$ alkynyl, $C_{3\text{-}6}$ cycloalkyl, hydroxy $C_{1\text{-}6}$ alkyl and R^3 is optionally mono, di- or tri-fluorinated, R4 and R5 are each independently hydrogen, $C_{1\text{-}6}$ alkyl, $C_{2\text{-}6}$ alkenyl, $C_{2\text{-}6}$ alkynyl, $C_{3\text{-}6}$ cycloalkyl or CF_3 or R^4 and R5, together with the nitrogen atom to which they are attached, form a 4-7 membered heteroaliphatic ring containing the nitrogen atom as the sole heteroatom, and R9 is benzyl or an aromatic ring containing either 6 atoms, 1, 2 or 3 of which are optionally nitrogen, or 5 atoms, 1, 2 or 3 of which are independently chosen from oxygen, nitrogen and sulphur, at most one of the atoms being oxygen or sulphur, and R9 is optionally substituted by one, two or three substituents independently chosen from halogen atoms and C1-4alkyl, C2-4alkenyl, C2-4alkynyl, C1-4alkoxy, $C_{2\text{-4}}$ alkenyloxy and $C_{2\text{-4}}$ alkynyloxy groups each of which groups is

WO 99/47171 PCT/GB99/00801

5

10

15

20

25

30

unsubstituted or substituted by one, two or three halogen atoms, and when X is a pyridine derivative, the pyridine derivative is optionally in the form of the N-oxide and providing that when X is a tetrazole derivative it is protected by a C_{1-4} alkyl group; or X is phenyl optionally substituted by one, two or three groups independently selected from halogen, cyano, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl and C_{3-6} cycloalkyl;

Y is optionally branched C_{1-4} alkylidene optionally substituted by an oxo group or Y is a group $(CH_2)_jO$ wherein the oxygen atom is nearest the group X and j is 2, 3 or 4;

Z is a 5-membered heteroaromatic ring containing 1, 2 or 3 heteroatoms independently selected from oxygen, nitrogen and sulphur, at most one of the heteroatoms being oxygen or sulphur and providing that when two of the heteroatoms are nitrogen an oxygen or sulphur atom is also present and that when one of the atoms is oxygen or sulphur then at least one nitrogen atom is present, or a 6-membered heteroaromatic ring containing 2 or 3 nitrogen atoms, Z being optionally substituted by Rv and/or Rw, where Rv is halogen, R6, NR7R8, NR7COR8, CN, furyl, thienyl, phenyl, benzyl, pyridyl or a 5-membered heteroaromatic ring containing at least one nitrogen atom and optionally 1, 2 or 3 other heteroatoms independently selected from oxygen, nitrogen and sulphur, at most one of the other heteroatoms being oxygen or sulphur and Rw is R6 or CN;

 R^6 is C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyloxy, C_{2-6} alkynyloxy, C_{1-6} alkynyloxy, C_{1-6} alkoxy, C_{2-6} alkynyloxy, C_{2-6} alkynyl

 R^7 and R^8 are each independently hydrogen, $C_{1\text{-}6}$ alkyl, $C_{2\text{-}6}$ alkenyl, $C_{2\text{-}6}$ alkynyl, $C_{3\text{-}6}$ cycloalkyl or CF_3 or R^7 and R^8 , together with the nitrogen atom to which they are attached, form a 4-7 membered heteroaliphatic ring containing the nitrogen atom as the sole heteroatom;

or a pharmaceutically acceptable salt thereof.

As used herein, the expression "C₁₋₆alkyl" includes methyl and ethyl groups, and straight-chained and branched propyl, butyl, pentyl and hexyl

groups. Particular alkyl groups are methyl, ethyl, n-propyl, isopropyl and t-butyl. Derived expressions such as " C_{1-4} alkyl", " C_{2-4} alkenyl", " C_{2-6} alkenyl", "hydroxy C_{1-6} alkyl", " C_{2-4} alkyl" and " C_{2-6} alkynyl" are to be construed in an analogous manner.

The expression "C₃₋₆cycloalkyl" as used herein includes cyclic propyl, butyl, pentyl and hexyl groups such as cyclopropyl and cyclohexyl.

5

10

15

20

25

Suitable 5- and 6-membered heteroaromatic rings include pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, oxadiazolyl and thiadiazolyl groups. A suitable 5-membered heteroaromatic ring containing four nitrogen atoms is tetrazolyl. Suitable 6-membered heteroaromatic rings containing three nitrogen atoms include 1,2,4-triazine and 1,3,5-triazine.

The term "halogen" as used herein includes fluorine, chlorine, bromine and iodine, of which fluorine and chlorine are preferred.

As used herein the term "C₁₋₆alkoxy" includes methoxy and ethoxy groups, and straight-chained, branched and cyclic propoxy, butoxy, pentoxy and hexoxy groups, including cyclopropylmethoxy. Derived expressions such as "C₂₋₆alkenyloxy", "C₂₋₆alkynyloxy", "C₁₋₄alkoxy", "C₂₋₄alkenyloxy" and "C₂₋₄alkyloxy" should be construed in an analogous manner.

Four particular compounds which can be used are: 6-(1-methylimidazol-4-yl)methyloxy-3-(5-methylisoxazol-3-yl)-1,2,4-triazolo[3,4-a]phthalazine;

3-(5-methylisoxazol-3-yl)-6-(1-methyl-1,2,3-triazol-4-yl)methyloxy-1,2,4-triazolo[3,4-a]phthalazine;

3-(5-methylisoxazol-3-yl)-6-(2-pyridyl)-1,2,4-triazolo[3,4-a]phthalazine; and

3-(5-methylisoxazol-3-yl)-6-(1-methylimidazol-4-yl)-1,2,4-triazol-3-ylmethyloxy-1,2,4-triazolo[3,4-a]phthalazine.

30 The second of the above compounds is particularly favoured.

The present invention also provides a pharmaceutical composition comprising an muscarinic agonist, an inverse agonist of the GABAA α_5 receptor subtype and a pharmaceutically acceptable carrier.

There is also provided a kit of parts comprising a first pharmaceutical composition comprising an muscarinic agonist and a first pharmaceutically acceptable carrier and a second pharmaceutical composition comprising an inverse agonist of the GABAA α_5 receptor subtype and a second pharmaceutically acceptable carrier for simultaneous, sequential or separate administration.

5

10

15

20

25

30

There is further provided a combination of an muscarinic agonist and an inverse agonist of the GABAA α_5 receptor subtype for use in a method of treatment of the human body, particularly for the treatment of a neurodegenerative disorder with associated cognitive deficit such as Alzheimer's Disease or Parkinson's disease, or of a cognitive deficit arising from a normal process such as aging or of an abnormal process such as injury. The combination is particularly beneficial in the treatment of Alzheimer's Disease.

There is also provided the use of a combination of an muscarinic agonist and an inverse agonist of the GABAA α_5 receptor subtype in the manufacture of a medicament for the treatment of a neurodegenerative disorder such as Alzheimer's Disease or Parkinson's disease, or of a cognitive deficit arising from a normal process such as aging or of an abnormal process such as injury. The treatment of Alzheimer's Disease is particularly preferred.

There is also disclosed a method of treatment of a subject suffering from a neurodegenerative disorder, such as Alzheimer's Disease or Parkinson's disease, or a cognitive deficit arising from a normal process such as aging or an abnormal process such as injury, which comprises administering to that subject a therapeutically effective amount of a combination of an muscarinic agonist and an inverse agonist of the GABAA

 α_5 receptor subtype. The treatment of Alzheimer's Disease is particularly preferred.

5

10

15

20

25

30

The pharmaceutical compositions of the present invention are preferably in unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, transdermal patches, auto-injector devices or suppositories; for oral, parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums or surfactants such as sorbitan monooleate, polyethylene glycol, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of each active ingredient of the present invention. Typical unit dosage forms contain from 1 to 100 mg, for example 1, 2, 5, 10, 25, 50 or 100 mg, of each active ingredient. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the

5

10

15

20

25

duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

For the treatment of a neurodegenerative condition, a suitable dosage level is about 0.01 to 250 mg/kg per day, preferably about 0.01 to 100 mg/kg per day, and especially about 0.01 to 5 mg/kg of body weight per day of each active ingredient. The compounds may be administered on a regimen of 1 to 4 times per day. In some cases, however, dosage outside these limits may be used.

The synergistic effect of the combination of the present invention can be shown, for example, by comparing the combined dosage of the combination with dosages of the same amount of each of the active ingredients separately on subjects using the Mini-Mental State Examination (MMSE) as described in Folstein and Folstein J. Psychiat. Res., 1975, 12, 189-198 or a variant thereof as discussed in Tombaugh and

McIntyre, JAGS, 1992, 40, 922-935.

CLAIMS

- A combination of an muscarinic agonist and an inverse agonist of the GABA_A α5 receptor subtype for separate, sequential or simultaneous aministration.
 - 2. A combination according to claim 1 wherein the inverse agonist has a functional efficacy at the α_5 receptor subtype of less than 20%, and a functional efficacy at the α_1 , α_2 and α_3 receptor subtypes of between -20 and +20 %.
 - 3. A combination according to claim 1 or 2 wherein the inverse agonist has a binding ration of greater than 10:1 to GABA_A receptors containing the α_5 receptor subtype compared to GABA_A receptors containing the α_1 , α_2 or α_3 subtypes.
 - 4. A combination according to claim 1 wherein the inverse agonist is 3-(5-methylisoxazol-3-yl)-6-(1-methyl-1,2,3-triazol-4-yl)methyloxy-1,2,4-triazolo[3,4-a]phthalazine.
 - 5. A combination according to any one of the preceding claims wherein the muscarinic agonist is selected from methacoline and its chloride, carbachol, bethanechol, arecholine, pilocarpine, muscarine, McN-A-343, oxetremorine, milameline, xanomeline, cis-methyldioxalene, pirenzepine, gallamine, SB 202026, AF102B, AFDX 116 and RS-86.
 - 6. A pharmaceutical composition comprising a combination as defined if any one of claims 1 to 5 and a pharmaceutically acceptable carrier for simultaneous administration.

25

10

15

20

7. A kit of parts comprising a first pharmaceutical composition comprising an muscarinic agonist and a first pharmaceutically acceptable carrier and a second pharmaceutical composition comprising an inverse agonist of the GABAA α_5 receptor subtype and a second pharmaceutically acceptable carrier for simultaneous, separate or sequential administration.

5

10

8. A method of treatment of a subject suffering from a neurodegenerative disorder or a cognitive deficit comprising administering to that subject a therapeutically effective amount of a combination of an muscarinic agonist and an inverse agonist of the GABAA $\alpha 5$ receptor subtype.

INTERNATIONAL SEARCH REPORT

International Application No PCT/GB 99/00801

. 2	OUT ON OF CUR LECT MATTER		
A. CLASSIFI	CATION OF SUBJECT MATTER A61K45/06		
According to	International Patent Classification (IPC) or to both national classification	n and IPC	
B. FIELDS S			
Minimum doc	cumentation searched (classification system tollowed by classification s	symbols)	
IPC 6	A61K		
Documentation	on searched other than minimum documentation to the extent that such	documents are included in the fields sea	rcnea
Electronic da	ata base consulted during the international search (name of data base	and, where practical, search terms used)	İ
C DOCUME	ENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant	ant passages	Relevant to claim No.
A	WO 98 04560 A (MERCK SHARP & DOHME	;REEVE	1-8
, ·	AUSTIN JOHN (GB); STERNFIELD FRANC	INE ()	
	5 February 1998 (1998-02-05)		
	abstract; claim 8		
A	WO 96 25948 A (MERCK SHARP & DOHME	;DAWSON	1-8
	GERARD RAPHAEL (GB))		
[29 August 1996 (1996-08-29) cited in the application		
1	abstract; claims		
			1 0
Α	WO 92 03433 A (NOVONORDISK AS)		1-8
	5 March 1992 (1992-03-05) page 1, line 34 - page 2, line 7		
	-	/	
[V] 5	rther documents are listed in the continuation of box C.	X Patent family members are listed	in annex.
		"T" later document published after the int or priority date and not in conflict with	the application but
"A" docur	nent defining the general state of the art which is not sidered to be of particular relevance	cited to understand the principle or to invention	neory underlying the
"E" earlie	r document but published on or after the international g date	"X" document of particular relevance; the cannot be considered novel or cannot be considered nov	or be considered to
"I " docur	nent which may throw doubts on priority claim(s) or th is cited to establish the publication date of another	involve an inventive step when the d	claimed invention
citat	ion or other special reason (as specified) ment referring to an oral disclosure, use, exhibition or	cannot be considered to involve an I	nventive step when the nore other such docu-
othe	er means	ments, such combination being obvi in the art.	ous to a person skilled
"P" docu late	ment published prior to the international filing date but r than the priority date claimed	"&" document member of the same pater	
Date of th	ne actual completion of the international search	Date of mailing of the international s	earch report
	24 August 1000	31/08/1999	
	24 August 1999		
Name an	nd mailing address of the ISA European Patent Office, P.B. 5818 Patentiaan 2	Authorized officer	
	European Patent Office, P.S. 3616 Fater Identical 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,	Leherte, C	
	Fax: (+31-70) 340-2040, 1x. 31 631 600 111,	Lenerte, C	

1

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 99/00801

	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	Relevant to claim No.
Category *	Citation of document, with indication where appropriate, of the relevant passages	
A	US 4 647 580 A (ROSZKOWSKI ADOLPH P) 3 March 1987 (1987-03-03) column 1, line 4,5 column 2, line 4-14	1-8

1

International application No.

PCT/GB 99/00801

INTERNATIONAL SEARCH REPORT

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
Claims Nos.: because they relate to subject matter not required to be searched by this Authority. namely:
2. X Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically: See FURTHER INFORMATION PCT/ISA/210
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This International Searching Authority found multiple inventions in this international application, as follows:
As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark on Protest The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claims relate to an extremely large number of possible combinations. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compositions claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those compounds explicitly mentionned in the claims

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/GB 99/00801

Patent document cited in search report		Publication date		atent family nember(s)	Publication date
WO 9804560	Α	05-02-1998	AU	3551997 A	20-02-1998
			AU	3553997 A	20-02-1998
			EP	0915877 A	19-05-1999
			EP	0915875 A	19-05-1999
			WO	9804559 A	05-02-1998
			NO	990304 A	25-03-1999
			ZA	9706591 A	18-08-1998
WO 9625948	 А	29-08-1996	AU	706515 B	17-06-1999
110 30203 10			AU	4725796 A	11-09-1996
			CA	2212058 A	29-08-1996
			EP	0810879 A	10-12-1997
			JP	11501302 T	02-02-1999
WO 9203433	 A	05-03-1992	AT	174027 T	15-12-1998
NO 3200 (00			AU	662105 B	24-08-1995
			AU	8403691 A	17-03-1992
			CA	2089769 A	22-02-1992
			CZ	9300244 A	19-01-1994
			DE	69130570 D	14-01-1999
			DE	69130570 T	29-04-1999
			EP	0544779 A	09-06-1993
			ES	2124707 T	16-02-1999
			FI	930747 A	30-03-1993
			HU	66863 A	30-01-1995
			HU	9500608 A	28-12-1995
			IL	99165 A	20-11-1997
			JP	6500542 T	20-01-1994
			NO	301883 B	22-12-1997
			NZ	239450 A	22-12-1994
			PT	98736 A,B	31-07-1992
			SK	12793 A	09-09-1993
			US	5418240 A	23-05-1995
			ÜS	5527813 A	18-06-1996
			ÜŠ	5578602 A	26-11-1996
			ÜS	5260314 A	09-11-1993
US 4647580	Α	03-03-1987	NON	 E	